

URINARY INCONTINENCE

Patient Fact Sheet

Introduction

Urinary incontinence is defined as the uncontrolled or involuntary loss of urine of any amount. Urinary incontinence affects as many as 25 million Americans, both men and women. Incontinence is a symptom, not a disease and is never due to “just old age.” Incontinence is a treatable and often curable condition.

There are four different types of incontinence:

Stress incontinence is the leakage of urine due to increased abdominal pressure when sneezing, laughing, lifting, changing position or exercising (i.e. jogging).

Urge incontinence is the leakage of urine due to a strong urge to urinate, usually on the way to the bathroom.

Mixed incontinence is a combination of both stress and urge incontinence.

Chronic retention of urine / Overflow Incontinence (a flaccid or enlarged bladder) is the leakage that occurs when the bladder cannot fully empty. It may be due to obstruction or injury such as enlarged prostate in men, scarring or narrowing of the urethra (as a result of previous surgery, congenital defects, sexually transmitted diseases or straddle injuries). Other causes include diabetes, stroke, multiple sclerosis, spinal cord injury and other neurological disease. Overflow incontinence may also be a side effect of medications.

Symptoms that you may notice:

- Leakage of urine with physical activities such as, exercise, coughing, laughing, sneezing or changing positions
- A strong urge to urinate followed by uncontrollable leakage
- A need to strain when passing urine
- Frequent urination; urinating more than 8 times per day or more than 2 times per night

Signs that your health care provider may find on examination:

- Leakage of urine when asked to cough or bear down during an examination with or without a full bladder
- Uncontrollable bladder contractions found on specialized bladder testing
- Large amounts of urine left in the bladder after voiding found on an ultrasound evaluation or other specialized testing
- Narrowing or blockage of the urethra found after specialized testing

Treatment:

- Limit your consumption of bladder irritants: alcohol, caffeine, artificial sweeteners, carbonated beverages, high sugar and citric acid foods and beverages.
- Retrain your bladder with regular, timed voiding throughout the day. When voiding, try to empty your bladder completely.
- Perform daily exercises to strengthen the pelvic muscles. Studies have shown that the best way to learn to do pelvic floor exercises is to work with a trained healthcare professional. To locate these muscles, try to stop the flow of urine during voiding (do not do this often) Perform quick contractions and long

contractions (holding and relaxing). You may do these exercises lying / sitting down or standing throughout the day.

- Discuss with your health care professional whether or not medications that can reduce uncontrollable urges to urinate are right for you. Also, ask about the possibility that your current medications may be affecting your bladder.
- Ask your health care professional if you need a referral to a specialist (Urologist, Gynecologist or Urogynecologist) to talk further about your diagnosis and treatment options.
- Treatment for stress incontinence include: lifestyle changes: weight loss, reducing dietary irritants, avoiding constipation, smoking cessation, pelvic floor muscle exercises, pessary placement in women and the injection of bulking agent such as collagen. Surgical treatments include: placement of an artificial urinary sphincter, bladder neck suspension or urethral sling.
- Treatment for urge incontinence include: fluid management; time and amount of fluid, limiting alcoholic and caffeine beverages, dietary management and weight reduction, avoiding constipation, pelvic floor exercises, bladder retraining and medications. Surgical treatments include: placements of sacral nerve stimulators and denervation surgery.
- Treatment for mixed incontinence includes: combinations of the above treatments.
- Treatment for overflow incontinence include: surgical repair of urethral strictures, removal or reduction of the obstruction, scheduled timed voiding, clean intermittent catheterization, and most importantly, addressing or curing the underlying condition or cause.

Prevention:

- Maintain a healthy weight. Obesity can lead to incontinence.
- Empty your bladder regularly (at least every 2-4 hours). It is also important to sit or stand in front of the toilet and wait for your bladder to empty completely.
- Practice pelvic muscle exercises regularly to strengthen the muscles that support your pelvic organs.
- Stop smoking. Coughing due to smoking can increase abdominal pressure and may contribute to stress incontinence. Nicotine may cause frequency and urgency leading to urge incontinence.
- Limit the use of alcohol. Alcohol can cause urgency and frequency leading to incontinence.
- Have a regular bowel routine. Constipation can lead to difficulty emptying the bladder. Maintaining an adequate amount of fiber, whole grains, fruits and vegetables in your diet will promote regularity.
- If you suspect a problem, keep a record of your diet and voiding habits and take them with you to your next appointment. This information will help your health care professional discuss any concerns you may have.
- Talk to your health care professional and find out how your medications may be affecting your bladder control. Changes in dosage or times may prevent incontinence.
- Be informed. Educate yourself by attending lectures, seminars and health fairs in your community to learn more about the causes of and treatment options for incontinence.

continued on back

Bladder Health Awareness Task Force

Susan Lipsy, MS, RN, CUNP, Chair

Pat Meade-D'Alisera, MSN, FNP, CURN, Co- Chair

Pamela Ceo MSN, APRN BC, CUNP

Cheryle Gartley, Simon Foundation

Mary Beth Hendricks, MSN, RNCS, FNP, ACNP, CUNP

Cheryl Le Croy, MSN, RN

Rachel Levkowicz, Health Educator NAFC

Silvia Maxwell, MSN, APRN BC, CUNP

Todd Morningstar, LPN

Donna Thompson, MSN, CRNP, CCCN

This material is for educational purposes only and should in no way be taken to be the practice or provision of medical, nursing or professional healthcare advice or services. The information should not be used in place of a visit, call, consultation or advice of your physician, nurse or other health care provider. The information obtained herein is not exhaustive and does not cover all aspects of the specific disease, ailment, physical condition or their treatments. Should you have any health care related questions, please call or see your physician, nurse or other health care provider promptly.

The Society of Urologic Nurses and Associates is a professional organization committed to excellence in clinical practice and research through education of its members, patients, family, and community.

© 2006 Society of Urologic Nurses and Associates



For more information, contact:

Society of Urologic Nurses and Associates

East Holly Avenue Box 56

Pitman, NJ 08071-0056

Phone 888-TAP-SUNA or 856-256-2335

suna@ajj.com

www.suna.org